



Contents

Purpose and procedure	66	Oestrogen and progestin increased urge and stress incontinence in postmenopausal women with coronary heart disease	84
EBN notebook			
The GATE frame: critical appraisal with pictures	68	A nurse led continence service reduced symptoms of incontinence, frequency, urgency, and nocturia.....	85
Become a sentinel reader for MORE EBN(McMaster Online Rating of Evidence)	72	Adding topical pale sulfonated shale oil to compression therapy and moist wound care reduced venous leg ulcer size but had no effect on complete wound healing after 20 weeks.....	86
Treatment			
Review: insufficient evidence exists on the effectiveness of proton pump inhibitors for adults with prolonged non-specific cough associated with gastro-oesophageal reflux	73	Review: sedative hypnotics may improve sleep quality but increase adverse effects in elderly people with insomnia.....	87
Review: some barrier creams and moisturisers are effective for irritant contact dermatitis; steroids are effective for allergic contact dermatitis	74	Review: protein and energy supplements reduce mortality and complications in undernourished elderly people in hospital	88
Review: lifestyle education programmes lower glucose concentrations and reduce the incidence of type 2 diabetes.....	75	A discharge planning intervention improved outcomes in older people admitted to hospital for hip fracture after a fall	89
Review: aspirin was effective for primary prevention of stroke in women and MI in men but increased major bleeding	76	Moving elderly inpatients to a transitional care facility reduced hospital stay but increased time to transfer to long term care.....	90
Review: secondary prevention programmes with and without exercise reduced all cause mortality and recurrent myocardial infarction.....	77	Clinical prediction guide	
Review: lorazepam provides the best control for status epilepticus	78	The STRATIFY falls risk assessment tool was not useful in predicting falls in patients with acute stroke.....	91
Review: laxatives in the form of fibre reduce symptoms in patients with symptomatic haemorrhoids	79	Qualitative	
Review: ginger prevents 24 hour postoperative nausea and vomiting	80	Personal beliefs, experiences, and emotions influenced smokers' perceptions of their cancer risk.....	92
Review: routine changes of IV administration sets (not containing lipids or blood products) at intervals \leq 96 hours do not affect infusate or catheter related bloodstream infection	81	Thresholds for requesting transfers to tertiary care were influenced by perceptions of the current illness and differences in perceptions of the initial hospital and tertiary care centre	93
Mechanical ventilation with heated humidifiers or with heat and moisture exchangers with microbiological filters did not reduce ventilator associated pneumonia in adults	82	Telephone support met the perceived needs of dementia caregivers for convenient access to information, referral, and emotional support	94
A pre-dialysis psychoeducational intervention increased length of survival more than usual care in patients with chronic kidney disease	83	Patients and their primary care providers identified 4 goals of life sustaining treatment.....	95
		Inappropriate use of medicines in acute care for the elderly related to a focus on acute care, providers' passive attitudes about learning, and paternalistic decision making	96
		Glossary	Inside back cover

Search all issues of *Evidence-Based Nursing* published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 47892.