



Contents

Purpose and procedure	2	Glargine dose titration by patients and by physicians were equally effective for preventing severe hypoglycaemia.....	20
EBN notebook			
Helping practitioners understand the contribution of qualitative research to evidence-based practice	4	Web based care management improved glucose control in patients with poorly controlled diabetes.....	21
Reflections on "Helping practitioners understand the contribution of qualitative research to evidence-based practice"	7	An educational booklet did not improve adherence or symptoms in patients prescribed thyroxine for primary hypothyroidism.....	22
Treatment			
The WHO technique for intramuscular thigh vaccination in infants and toddlers had fewer adverse reactions than 2 other techniques	9	Review: multidisciplinary interventions reduce hospital admission and all cause mortality in heart failure.....	23
Liposomal lidocaine improved intravenous cannulation success rates in children	10	Review: multivitamins and mineral supplements do not reduce infections in elderly people	24
Review: children permitted clear fluids \leq 120 minutes before surgery have similar gastric volumes and pH values as those on standard fasts.....	11	Multivitamin and multimineral supplements did not reduce reported infection days or related use of healthcare services in elderly people.....	25
Topical chloramphenicol was not effective for acute infective conjunctivitis in children.....	12	Clinical prediction guide	
Nurse home visits did not differ from standard care for prevention of recurrent child abuse	13	A simple risk score predicted 7 day stroke risk after transient ischaemic attack	26
Multisystemic therapy improved adherence to blood glucose testing in adolescents with type 1 diabetes....	14	Qualitative	
A weight maintenance diet reduced bulimic symptoms in adolescent girls.....	15	Mothers' decisions about MMR vaccination were framed by their children's vulnerabilities and wider social trends.....	27
After 3 months, low dose oral contraceptives reduced pain in adolescent girls with moderate to severe dysmenorrhoea	16	UK and US adolescents perceived internet health information to be salient but of questionable credibility	28
Trimethoprim reduced dysuria in women with symptoms of urinary tract infection but negative urine dipstick test results.....	17	Young women described the benefits of having advance supplies of emergency contraception but emphasised its use as a "last resort" rather than an alternative form of contraception.....	29
Review: wearing graduated compression stockings during air travel reduces the risk of deep venous thromboembolism.....	18	Marginalised HIV positive drug users felt that an HIV diagnosis created benefits as well as losses	30
A home based, physical activity intervention increased physical activity, fitness, and vigour and reduced fatigue in sedentary women with early stage breast cancer ...	19	Perceived barriers and benefits were factors in decision making about colorectal screening.....	31
		UK palliative care professionals identified service infrastructure, patient and carer attitudes and characteristics, and practice culture as influencing place of death of patients with cancer.....	32
		Glossary	Inside back cover

Search all issues of *Evidence-Based Nursing* published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 13845.