

# GERIATRIC NURSING

VOL. 38, NO. 2 □ MARCH/APRIL 2017

## EDITORIAL

### FROM THE EDITOR

- 95 From Capstone/Dissertation to publication: Let's try this again!**

Barbara Resnick

## FEATURE ARTICLES

- 97 Effectiveness of a community-based program for suicide prevention among elders with early-stage dementia: A controlled observational study**

Jong-Pill Kim and Jinhyang Yang

This community-based program in geriatric nursing practice can be effective in preventing suicide among elders with early stage dementia.

- 106 Compassion fatigue among nurses working with older adults**

Kay L. Kolthoff and Susan E. Hickman

The study suggests that more experienced nurses have less compassion fatigue in comparison to newer nurses. It is possible that nurses who are better able to cope stay in their positions longer than less experienced nurses.

- 110 Caregiver's distress related to the patient's neuropsychiatric symptoms as a function of the care-setting**

Laura Lorenzo-López,  
Carmen de Labra, Ana Maseda,  
Trinidad Lorenzo, Helena Agrafojo,  
José Luis Rodríguez-Villamil,  
Isabel González-Abraldes, and  
José C. Millán-Calenti

While high patients' scores in hallucinations, sleep and night-time disturbances, anxiety, and aberrant motor behavior were the best predictors of professional caregiver's distress, agitation/aggression, delusions, disinhibition, apathy/indifference, depression/dysphoria, and elation/euphoria were the best predictors of informal caregiver's distress.

- 119 Development and psychometric testing of a measure of older adult patients' attitudes towards mobility during hospitalization (ATM-H)**

Chedva Levin, Nurit Gur-Yaish,  
Maayan Agmon, Efrat Gil,  
Debbie Rand, Barbara King, and  
Anna Zisberg

The aim of this study is to develop a measure of patients' attitudes towards mobility in the hospital setting and to test its psychometric properties.

- 124 Trajectories of health-related quality of life among family caregivers of individuals with dementia: A home-based caregiver-training program matters**

Li-Min Kuo, Huei-Ling Huang,  
Jersey Liang, Yam-Ting Kwok,  
Wen-Chuin Hsu, Chin-Yi Liu,  
and Yea-Ing L. Shyu

Caregivers who received the training program were more likely than those who did not to have health-related quality of life that followed a well-functioning trajectory over 18 months.