

Departments



6 EDITOR'S NOTE

Should you go to jail if you make an

8 LETTERS

Put your best foot forward for a choking victim...reporting a colleague for substance abuse...why Mcn in Nursing?

10 ADVICE, P.R.N.

Slacker on the unit...full name required on ID badge?...tetanus prophylaxis

12 MEDICATION ERRORS

Infusing promethazine safely, sorting out soundalike drug names

14 PATIENT SAFETY

Insulin safety in the hospital

18 WOUND AND SKIN CARE

Measuring wounds

22 INSIGHTS ON DEATH AND DYING

Permission to mourn...avoiding deception...pain management



24 CLINICAL QUERIES

Responding to a chronically ill patient who's suicidal

26 CONTROLLING PAIN

New options: Techniques and devices

28 COMBATING INFECTION

Stamping out scabies

31 DRUG NEWS

Deaths linked to use of methadone for analgesia...pharmacy program enhances adherence to drug regimens...two bone-huilding drugs compared... pregabalin eases pain of fibromyalgia... anti-malarial drug regains effectiveness

66 I.V. ROUNDS

Managing hyperemesis gravidarum

69 CLINICAL DO'S AND DON'TS

Assessing a patient for abuse

70 EYE ON DIAGNOSTICS

Screening for microalbuminuria

71 DRUG CHALLENGE

Match drug names to their ingredients and indications



80 ACTION STAT Intussusception

By Heather Brown-Guttovz, RN, BSN Act fast to prevent intestinal necrosis, sepsis, and death.

17 About Nursing2007

- 72 Advertisers' Index
- 74 Classified Advertisements

*After page 64...

Hospital section feature:

Community-acquired pneumonia

Critical care feature:

Holiday heart syndrome

*If you work in a hospital or in critical care but don't see that section in this issue, please call our customer service department at 1-800-879-0498.

Practice pointers from this issue of Nursing2007

- About the size of a credit card, the new IONYSYS patient-controlled transdermal system delivers 80 doses of fentanyl and requires no pump (ρ. 27).
- **Aortic aneurysms** are commonly linked to atherosclerosis, but they can also develop in people with "clean" arteries (p. 38).
- Change the wound treatment regimen if a wound's measurement hasn't decreased in 2 weeks (p. 18).
- When giving first aid for a cut or puncture of the eye or eyelid, don't wash the eye with water or any other liquid (p. 47).
- A diagnosis of fibromyalgia takes an average of 5 years because of the lack of diagnostic tests specific to the disorder (p. 63).
- The scabes mite can't live for more than 3 days without a human host and isn't spread by casual contact (p. 28).

CEs coming next month

Sepsis

Managing pain in patients who are opioid-tolerant Lower extremity ulcers

On the Web...

Go to www.Nursing2007.com to take the nurse/physician relationship survey.